

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



B1

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶:
C12N 15/00, 15/05, 15/09, 15/11, 15/29,
15/64, 15/82, A01H 1/00, 3/00, 4/00

A1

(11) International Publication Number: WO 97/25419

(43) International Publication Date: 17 July 1997 (17.07.97)

(21) International Application Number: PCT/US97/00510

(22) International Filing Date: 10 January 1997 (10.01.97)

(30) Priority Data:
08/586,331 12 January 1996 (12.01.96) US

(71) Applicant: THE UNITED STATES OF AMERICA, represented by TH [US/US]; E SECRETARY OF AGRICULTURE, Washington, DC 20250 (US).

(72) Inventors: BLECHL, Ann, E.; 1005 Buchanan Street, Albany, CA 94706 (US). ANDERSON, Olin, D.; 208 Poshard Street, Pleasant Hill, CA 94523 (US).

(74) Agents: PENDORF, Stephan, A. et al.; Suite 1000, 600 N. Westshore Boulevard, Tampa, FL 33609 (US).


(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published


With international search report.

(54) Title: GLUTENIN GENES AND THEIR USES

Y-type HMW-glutenin



X-type HMW-glutenin



(57) Abstract

The present invention is directed to methods of improving flour by altering glutenin content in seeds of plants, particularly wheat. The invention relates to methods of introducing a recombinant construct comprising a glutenin gene into a parental plant.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

GLUTENIN GENES AND THEIR USES

Field of the Invention

The present invention relates generally to plant molecular biology. In particular, it relates to nucleic acids and methods for improving glutenin content of plants.

BACKGROUND OF THE INVENTION

The glutenins, which include both high molecular weight (HMW) glutenin subunits and low molecular weight (LMW) glutenin subunits, comprise an economically important class of wheat seed storage proteins. The apparent molecular weights of the individual HMW glutenin polypeptides or subunits range from 90 to 200 kDa. These subunits crosslink by disulfide bonds among themselves and with LMW glutenin polypeptides to form polymers exceeding one million daltons in molecular weight. HMW glutenins constitute 8-10%, while LMW glutenins constitute 15-20% of the total endosperm protein. Both HMW and LMW glutenin proteins play important functional roles in determining the end-uses of wheat flour.

In wheat, HMW glutenins are encoded at the Glu-1 loci on the long arms of the group 1 chromosomes. Each locus consists of two separate genes, encoding an x-type and a y-type subunit, respectively. These pairs have never been confirmed to be separated by recombination. This has made determination of their separate contributions to bread dough properties difficult to assess by genetic correlation studies. For a review of the genetics and biochemistry of glutenin polypeptides, see, Shewry et al., J. Cereal Sci. 15:105-120 (1992).

Both the quantity and identity of specific HMW glutenin alleles contribute to the differences in bread-making quality of various cultivars. For instance, deletion of glutenin genes results in a decrease in the overall levels of HMW glutenins, which results in decreases in bread-making quality (see, e.g., Lawrence et al. J. Cereal Sci 7:109-112 (1988)).

The effects of overproducing HMW glutenin on protein accumulation and baking quality has not been assessed because such lines of wheat have not been found among natural

populations. In addition, direct alteration of the glutenin subunits that form the polymers is not possible using standard breeding methods. Thus, the art lacks reproducible and efficient methods of producing lines with altered glutenin contents. The present invention addresses these and other needs.

SUMMARY OF THE INVENTION

The present invention provides methods of increasing glutenin in the endosperm of wheat plants. The methods and plants of the invention therefore are useful in providing flour and dough having improved end-use properties. The methods comprise introducing into a parental wheat plant a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and selecting progeny wheat plant having increased glutenin content in the endosperm of mature seed. The glutenin content of the progeny is preferably at least about 15% greater than the glutenin of the parental wheat plant.

Any method of introducing the expression cassette into the parental plant can be used. Particle bombardment is a convenient method for producing transgenic plants. Once the expression cassette is stably integrated into the genome, standard sexual crosses can be used to introduce the expression cassette into desired lines.

In some embodiments, the nucleic acid introduced into the plant encodes a chimeric glutenin polypeptide. For instance, the chimeric polypeptide may comprise sequences from an x-type glutenin polypeptide and a y-type glutenin polypeptide. Exemplary genes for this purpose include the Glu-D1-1b gene and the Glu-D1-2b gene.

The expression cassette may further comprise a seed-specific promoter to direct expression of the introduced nucleic acid to the endosperm. A convenient promoter for this purpose is the promoter from the Glu-D1-2b gene.

Any wheat cultivar can be used as the parental line in the present invention. An exemplary cultivar is Bobwhit.

The invention also provides wheat plants comprising a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide. The plants of the invention

have a glutenin content in the endosperm of a mature seed at least about 15% greater than the glutenin content in the endosperm of a mature seed from a parental wheat plant. The percentage of total endosperm protein in which is glutenin will usually depend upon the glutenin content of the parental line. Usually, HMW glutenins account for at least about 15% of the total protein in the endosperm of the mature seed from plants of the invention.

The invention further provides recombinant constructs comprising a wheat glutenin gene promoter of about 400 to about 2800 nucleotides operably linked to a heterologous polynucleotide sequence. These constructs are particularly useful in directing expression of the heterologous sequence to seeds of transgenic plants. An exemplary wheat glutenin gene from which the promoters of the invention can be derived is Glu-D1-2b.

Definitions

The term "plant" includes whole plants, plant organs (e.g., leaves, stems, roots, flowers, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

A "heterologous sequence" is one that originates from a foreign species, or, if from the same species, is substantially modified from its original form. For example, a promoter operably linked to a heterologous structural gene is from a species different from that from which the structural gene was derived, or, if from the same species, one or both are substantially modified from their original form.

A "glutenin polypeptide" is a gene product of a glutenin gene or glutenin polynucleotide sequence. A glutenin polypeptide can be either a LMW glutenin or a HMW glutenin. A glutenin polypeptide contains cysteine residues by which disulfide bonds are formed with other glutenin polypeptides to form polymers. The composition and size of the repeat region is also important to polymer formation.

A "chimeric glutenin polypeptide" is glutenin gene product that comprises a modified amino acid sequence. Modifications, as explained in detail below, can be in the form of substitutions, deletions, or additions of single amino acids or groups of amino acids. Thus, chimeric glutenin polypeptides can be hybrid glutenin polypeptides comprising sequences from two or more different subunits or may be polypeptides in which single amino acid modifications are made.

In the case where an inserted polynucleotide sequence is transcribed and translated to produce a functional glutenin polypeptide, one of skill will recognize that because of codon degeneracy, a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the terms "glutenin gene" or "glutenin polynucleotide sequence". In addition, the terms specifically include those full length sequences substantially identical (determined as described below) with a glutenin gene sequence and that encode proteins that retain the function of the glutenin polypeptide. Thus, in the case of wheat glutenin genes disclosed here, the term includes variant polynucleotide sequences which have substantial identity with the sequences disclosed here and which encode glutenin polypeptides capable of crosslinking by disulfide bonds with other glutenin polypeptides to form glutenin polymers.

Two polynucleotides or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the complementary sequence is identical to all or a portion of a reference polynucleotide sequence.

Sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a segment or "comparison window" to identify and compare local regions of sequence similarity. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman Adv. Appl. Math. 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch J. Mol. Biol. 48:443 (1970), by the search for similarity method of Pearson and Lipman Proc. Natl.

Acad. Sci. (U.S.A.) 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection. "Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 60% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using the programs described above using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids

having aromatic side chains is phenylalanin , tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine , and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine , lysine-arginine, alanine-valine, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C to about 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. However, nucleic acids which do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram showing the domains of y-type and x-type HMW glutenins. The C- and N-terminal domains are shaded, the locations of cysteine residues are designated by S.

Figure 2A is a diagram showing the structure of a hybrid glutenin polypeptide of the invention.

Figure 2B is a diagram of the recombinant plasmid of the invention which encodes the polypeptide shown in Figure 2A.

Figure 3 is a photograph of a gel stained with Coomassie blue showing the relative mobility of a hybrid HMW glutenin polypeptide of the invention (marked by the arrow).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

This invention relates to plant glutenin genes, in particular wheat glutenin genes. Nucleic acid sequences from glutenin genes can be used to modify glutenin content in transgenic plants, in particular wheat plants. In addition,

th invention provides new seed-specific promoters useful for directing expression of desired genes in a number of plants.

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning - A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

Glutenin polypeptides and genes

The HMW glutenins are composed primarily of a central domain of 45-90 repeating two or three simple motifs, comprising chiefly glutamine and proline. This unusual primary structure results in a rod-like secondary structure as assessed by circular dichroism and SEM. Cysteines in the C- and N-terminal domains of the molecules are known to be critical for the formation of intermolecular disulfide bonds (Figure 1). The quaternary interactions of these proteins are of interest because the disulfide crosslinks among HMW and LMW subunits is thought to be integral to the elastic properties of dough.

Any isolated glutenin gene can be used in the present invention. The particular polynucleotide sequence used is not a critical feature of the invention, so long as the desired alteration in glutenin content is achieved. As noted above, HMW glutenins (both x-types and y-types) are encoded at the Glu-1 locus. In hexaploid wheat the group 1 chromosomes are designated 1A, 1B and 1D. Studies of the frequency of alleles at each of the loci indicate the presence of at least three alleles at Glu-A1, 11 alleles at Glu-B1, and six alleles at Glu-D1. The gene products of each allele are designated by noting the chromosome, followed by its classification as x or y, and a number (see, MacRitchie, Advances in Food and Nutrition Research 36:1-87 (1992)).

While HMW glutenin genes have been cloned and described in the literature. For instance, a complete set of HMW glutenin

genes have been isolated and sequenced from the hard red winter wheat at Cheyenne (Anderson et al. Wheat Genetics Symposium Proc. pp 699-704 (Bath Press, Cambridge, UK, 1988), including two, Dx5 and Dyl0, with the highest correlation with good flour quality (Anderson et al. Nuc. Acids. Res. 17:461-462 (1989)). Other particularly useful HMW glutenins include Ax and Bx alleles.

The isolation of other glutenin genes may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed in the prior art can be used to isolate the desired gene from a cDNA or genomic DNA library. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a cDNA library, mRNA is isolated from endosperm and a cDNA library which contains the glutenin gene transcript is prepared from the mRNA.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology to amplify the sequences of the glutenin and related genes directly from genomic DNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds., Academic Press, San Diego (1990)).

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers et al., Cold Spring Harbor Symp. Quant. Biol. 47:411-418 (1982), and Adams et al., J. Am. Chem. Soc. 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding

th complementary strand using DNA polymerase with an appropriate primer sequence.

Isolated sequences prepared as described herein can then be used to modify glutenin gene expression and therefore glutenin content in plants. One of skill will recognize that the nucleic acid encoding a functional glutenin protein need not have a sequence identical to the exemplified genes disclosed here. Thus, genes encoding chimeric glutenin polypeptides can be used in the present invention.

As noted above, glutenin polypeptides, like other proteins, have different domains which perform different functions. Thus, the glutenin gene sequences need not be full length, so long as the desired functional domains of the protein is expressed. Chimeric glutenin polypeptides can be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art. For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. In particular, cysteine residues may be added, deleted or moved within the polypeptide to achieve a modified glutenin polypeptide with desired properties. Chimeric polypeptides may also be produced by fusing coding sequences from two or more glutenin genes. All of these modifications can be used in a number of combinations to produce the final modified protein chain.

Preparation of recombinant constructs

To use isolated glutenin sequences in modifying glutenin content in plants, recombinant DNA vectors suitable for transformation of plant cells are prepared. A DNA sequence coding for the desired glutenin polypeptide, for example a cDNA or a genomic sequence encoding a full length protein, is conveniently used to construct a recombinant expression cassette which can be introduced into the desired plant. An expression cassette will typically comprise the glutenin polynucleotide sequence operably linked to a promoter sequence and other transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the glutenin gene in the intended tissues (e.g., endosperm) of the transformed plant.

For example, a constitutive plant promoter fragment may be employed which will direct expression of the glutenin in all tissues of a plant. Such promoters are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may be under environmental control. Such promoters are referred to here as "inducible" promoters. Examples of environmental conditions that may effect transcription by inducible promoters include pathogen attack, anaerobic conditions, or the presence of light.

Typically, the promoters used in the constructs of the invention will be "tissue-specific" and are under developmental control such that the desired gene is expressed only in certain tissues, such as leaves, roots, fruit, seeds, or flowers. Promoters that direct expression in seeds, particularly the endosperm are particularly preferred. Examples of such promoters include the promoter from genes encoding seed storage proteins, such as napin, cruciferin, phaseolin, and the like (see, U.S. Patent No. 5,420,034). Other promoters suitable for expressing glutenin genes in cereals include promoters from genes encoding gliadins, cereal prolamines (e.g., zein, hordein, secalin, and avenin) and starch biosynthetic enzymes.

The endogenous promoters from glutenin genes are particularly useful for directing expression of glutenin genes to the seed, particularly the endosperm. These seed-specific promoters can also be used to direct expression of heterologous structural genes. Thus, the promoters can be used in recombinant expression cassettes to drive expression of any gene whose expression in seeds is desirable. Examples include genes encoding proteins useful in increasing the nutritional value of seeds (e.g., genes encoding proteins involved in lipid, protein, and carbohydrate or starch biosynthesis). Other genes include those encoding pharmaceutically useful

compounds, and genes encoding plant resistance products to combat fungal or other infections of the seed.

The glutenin promoters can also be used to initiate transcription of mRNA molecules to inhibit expression of an endogenous endosperm gene. Exemplary genes whose expression could be inhibited include genes encoding glutenins, enzymes involved in carbohydrate or lipid synthesis, secalins, and the like. Means for inhibiting gene expression in plants using recombinant DNA techniques are well known. For instance, antisense technology can be conveniently used. see, e.g., Sheehy et al., Proc. Nat. Acad. Sci. USA, 85:8805-8809 (1988), and Hiatt et al., U.S. Patent No. 4,801,340. Catalytic RNA molecules or ribozymes can also be used to inhibit expression of endosperm-specific genes. The design and use of target RNA-specific ribozymes is described in Haseloff et al. Nature, 334:585-591 (1988). Introduction of nucleic acid configured in the sense orientation has also been shown to be an effective means by which to block the transcription of target genes. For an example of the use of sense suppression to modulate expression of endogenous genes see, Napoli et al., The Plant Cell 2:279-289 (1990), and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

The glutenin promoters of the invention are typically at least about 400 base pairs in length, and often at least about 800 or about 1000 base pairs. The length of the promoters is typically less than about 3500 base pairs, usually less than about 2800 base pairs and often less than about 2000 base pairs in length. The length of the promoters is counted upstream from the translation start codon of the native gene. One of skill will recognize that use of the "about" to refer to lengths of nucleic acid fragments is meant to include fragments of various lengths that do not vary significantly from the lengths recited here and still maintain the functions of the claimed promoters (i.e., seed-specific gene expression).

To identify glutenin promoters, the 5' portions of a genomic glutenin gene clone is analyzed for sequences characteristic of promoter sequences. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In plants, further upstream from the

TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing et al., in Genetic Engineering in Plants, pp. 221-227 (Kosage, Meredith and Hollaender, eds. 1983).

In preparing expression vectors of the invention, sequences other than the promoter and the structural gene of interest are also preferably used. If proper polypeptid expression is desired, a polyadenylation region at the 3'-end of the glutenin coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences from a glutenin gene will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron, or phosphinothricin (the active ingredient in bialaphos and Basta).

Preparation of transgenic plants

The DNA constructs described above may be introduced into the genome of the desired plant host by a variety of conventional techniques. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising et al., Ann. Rev. Genet. 22:421-477 (1988).

The DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as biolistic methods, electroporation, PEG poration, and microinjection of plant cell protoplasts or embryogenic callus. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced using an Agrobacterium tumefaciens or A. rhizogenes vector.

Particle bombardment techniques are described in Klein et al., Nature 327:70-73 (1987). A particularly preferred method of transforming wheat and other cereals is the bombardment of calli derived from immature embryos as described by Weeks et al. Plant Physiol. 102:1077-1084 (1993).

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA* 82:5824 (1985).

Agrobacterium tumefaciens-mediated transformation techniques are also well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983). Although *Agrobacterium* is useful primarily in dicots, certain monocots can be transformed by *Agrobacterium*. For instance, *Agrobacterium* transformation of rice is described by Hiei et al. *Plant J.* 6:271-282 (1994).

The present invention is particularly useful in wheat and other cereals. A number of methods of transforming cereals have been described in the literature. For instance, transformation of rice is described by Toriyama et al. *Bio/Technology* 6:1072-1074 (1988), Zhang et al. *Theor. Appl. Gen.* 76:835-840 (1988), and Shimamoto et al. *Nature* 338:274-276 (1989). Transgenic maize regenerants have been described by Fromm et al., *Bio/Technology* 8:833-839 (1990) and Gordon-Kamm et al., *Plant Cell* 2:603-618 (1990)). Similarly, oats (Sommers et al., *Bio/Technology* 10:1589-1594 (1992)), wheat (Vasil et al., *Bio/Technology* 10:667-674 (1992)); Weeks et al., *Plant Physiol.* 102:1077-1084 (1993)), sorghum (Casas et al., *Proc. Natl. Acad. Sci. USA* 90:11212-11216 (1993)), rice (Li et al., *Plant Cell Rep.* 12:250-255 (1993)), barley (Yuechun and Lemaux, *Plant Physiol.* 104:37-48 (1994)), and rye (Castillo et al., *Bio/Technology* 12:1366-1371 (1994)) have been transformed via bombardment.

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocid and/or herbicid marker which has been introduced together with the glutathione polynucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation*

and Cultures, Handbook of Plant Cell Culture, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, Regeneration of Plants, Plant Protoplasts, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. Ann. Rev. of Plant Phys. 38:467-486 (1987).

The methods of the present invention are particularly useful for incorporating the glutenin polynucleotides into transformed plants in ways and under circumstances which are not found naturally. In particular, the glutenin polypeptides may be expressed at times or in quantities which are not characteristic of natural plants.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

The invention has use over a broad range of types of plants. The glutenin genes are preferably expressed in cereal species commonly used for production of flour, e.g., wheat, rye, oats, and the like.

The seed-specific promoters from glutenin genes can be used in essentially any plant species. For instance, the promoters or genes described here can be used in species from the genera Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Cucumis, Cucurbita, Daucus, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Juglans, Lactuca, Linum, Lolium, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Oryza, Panicleum, Pannisetum, Persea, Phaseolus, Pisum, Pyrus, Prunus, Raphanus, Secale, Senecio, Sinapis, Solanum, Sorghum, Trigonella, Triticum, Vitis, Vigna, and, Zea. Plants that are particularly useful in the invention include Avena, Brassica, Glycine, Hordeum, Oryza, Phaseolus, Pisum, Secale, Sorghum, Triticum, Vigna, and Zea.

As noted above, the invention is particularly useful for improvement of wheat cultivars. Any wheat cultivar may be improved by the methods of the invention. Exemplary lines

include hexaploid lines such as Anza, Shasta, Yorko Rio, Siouxland, Freedom, Tam107, and Tam200.

The effect of the modification of glutenin gene expression can be measured by detection of increases or decreases in desired glutenin protein levels using, for instance, gel electrophoresis, as described below. Quantification of HMW glutenin content can be carried out by SDS/PAGE densitometry, as described below.

Other methods for quantifying glutenin content include sonication of flour dispersions in SDS buffer to solubilize glutenin and other relatively insoluble proteins (see, Singh and MacRitchie in *Wheat End-Use Properties: Wheat and Flour Characterization for Specific End-Uses* H. Salovaara, ed., pp 321-326 (University of Helsinki, Helsinki, 1989)). This method is useful in solubilizing at least 95% of total flour protein.

Quantification of the glutenin fraction can then be performed using size exclusion high performance liquid chromatography as described by Batey et al., *Cereal Chem.* 68:207-209 (1991). Other suitable methods include reverse phase HPLC and capillary electrophoresis.

The plants of the invention have increased in glutenin content in the endosperm of mature seeds as compared to the parental lines from which they are derived. The increase is usually at least about 15% as compared to the parental line. More usually, the increase is at least about 20%, preferably at least about 35%, and in more preferred embodiments at least about 45%. Any of the methods for quantifying glutenins in the endosperm described above can be used to determine the percent increase in glutenin content. A wheat seed typically reaches maturity about 6 to about 10 weeks after anthesis.

Another measure of increased glutenin content in transgenic plants of the invention is glutenin content as a percent of total protein of a mature seed or flour derived from the mature seed. Different wheat cultivars have different glutenin contents as measured using standard methods. Thus, the final glutenin content of the plants of the invention will depend upon the parental line. In the case where HMW glutenin genes are used, the plants of the invention will typically have HMW glutenin contents of at least about 15%, usually at least

about 18%, and preferably at least about 20% as measured using standard methods such as those described in detail below.

Analysis of the rheological properties of flour derived from the transgenic plants of the invention can be carried out according to standard physical dough-testing instruments widely used to measure flour and dough quality (see, e.g., MacRitchie, *Advances in Food and Nutrition Research* 36:1-87 (1992)). Such methods include, for instance, use of extensographs to measure tensile strength. Two of the main parameters measured are maximum resistance (R_{max}) and extensibility (Ext). Other methods include mixographs and bake-test loaf volume (MacRitchie and *Gras Cereal Chem.* 50:292-302 (1973) as well as SDS-sedimentation tests, amelographs, and cookie spread methods.

The following Examples are offered by way of illustration, not limitation.

Example 1

This example demonstrates increase in glutenin content in transgenic wheat plants comprising recombinant constructs of the invention.

MATERIALS AND METHODS

Construction of the hybrid HMW glutenin expression plasmid

Two HMW glutenin subunits, Dx5 and Dy10, encoded by the Glu-D1-1b and Glu-D1-2b genes, respectively, have been most often associated with superior dough strength by genetic correlation studies. A bacterial expression vector comprising a fusion of coding regions of the Dy10 and Dx5 genes is described by Shani et al., *Plant Physiol.* 98:433-441 (1992). The hybrid HMW glutenin polypeptide encoded by this gene is diagrammed in Figure 2A.

In this example, the Dy10:Dx5 hybrid coding region was reunited with 5' and 3' flanking sequences from the native Dy10 and Dx5 genes. The construction of a fusion at the HindIII site located at base pairs 429-435 and 385-391 after the A of the start codons of the Glu-D1-2b (Genbank Accession No. X12929) and Glu-D1-1b (Genbank Accession No. X12928) genes (Anderson et al. *Nuc. Acids Res.* 17:461-462 (1989)) and its

insertion into the bacterial expression vector pet3A to form p t-3a-10/5 is described by Shani et al., supra.

Thus, the native Dyl0 gene provides the promoter, transcription start site, the 5' transcribed untranslated region and the first 145 codons including the 21 amino acid signal peptide. The native Dx5 gene provides its codons 130 to 848 followed by two translation stop codons, the transcription termination and poly(A) addition signals. Because amino acids 138 through 147 are shared by the Dyl0 and Dx5 glutenin subunits, the central region of repeating amino acid motifs of the hybrid glutenin polypeptide (thickest boxes in Figure 2A) is identical to that of Dx5. The configuration of the cysteines (S) and thus of potential disulfide bonds is chimeric in nature: the N-terminal cysteines are of the y-type and the single C-terminal cysteine is typical of x-type subunits.

In order to reconstitute the promoter region, transcription start and termination sites, and any other regulatory signals needed for expression in wheat, three fragments were isolated and combined in a single ligation reaction with the Bluescript KS⁻ (Stratagene) vector cut with EcoRI and BamHI: 1) the 715 bp StuI/DrdI fragment that contains the junction region in pET-3a-10/5; 2) the 2800 bp fragment from a clone of the native Dyl0 gene beginning at the EcoRI site 5' to the gene and ending at the StuI site 74 bp after the A in the start codon; and 3) the 3800 bp fragment from a clone of the native Dx5 gene that starts at the DrdI site 744 bp after the A in the start codon and ends at a BamHI site in the vector just outside the EcoRI site in the 3' flanking region. A plasmid with the correct structure was identified by restriction analysis and named pGlul0H5 (Figure 2B). This plasmid was deposited under the terms of the Budapest Treaty in the Agricultural Research Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois, USA, on January 12, 1996 and has been assigned Accession No. NRRL B-21517.

The plasmid UBI:BAR which comprises the maize ubiquitin promoter operably linked to the bar gene encoding the enzyme PAT, which inactivates phosphinothricin was prepared as described Cornejo et al. Plant Mol. Biol. 23:567-581 (1993).

Both plasmids were prepared by the alkaline lysis method using a Qiagen kit for the final purification. The DNAs were stored in TE buffer at a concentration of 1 mg/ml.

Wheat Transformation

12.5 ug of each DNA (an approximately 2:1 molar ratio of UBI:BAR to pGlu10H5) was coated onto gold particles for bombardment into immature wheat embryos of cultivar Bobwhite essentially as described by Weeks et al. Plant Physiol. 102:1077-1084 (1993) except that the embryos were incubated on callus induction media containing mannitol for 4 hours before and 20 hours after bombardment. Transformants were selected on the basis of their resistance to 1mg/L bialaphos at the callus and green shoot stages of regeneration and to 3 mg/L bialaphos at the rooting stage of regeneration as described by Weeks et al., supra.

Protein Analysis

Protein extracts from immature wheat endosperm were prepared by grinding the tissue or cells in SDS-PAGE sample loading buffer [66 mM Tris pH 6.8, 3% (w/v) SDS, 0.05% (w/v) Bromphenol blue, 2% (v/v) β -mercaptoethanol]. Protein extracts from mature dry wheat seeds were prepared by dissolving flour in sample buffer. SDS-PAGE was according to standard procedures. The separation gels for detection and quantitation of the hybrid subunit were 10% (w/v) acrylamide 0.05% (w/v) bis-acrylamide. The gels were run at 40 mamps constant current until 20 minutes after the bromphenol blue dye front had run off the gel (about 1000 Volt-hours).

The gels were stained by the Neuhoff method with a colloidal suspension of Coomassie blue. Protein standards for molecular weight calibration were purchased from Pharmacia and consisted of Phosphorylase b (94 kDa) Bovine Serum Albumin (67 kDa), Ovalbumin (43 kDa), Carbonic Anhydrase (30 kDa), Soybean Trypsin Inhibitor (20.1 kDa) and α -Lactalbumin (14.4 kDa). Stained gels were scanned using an Alpha Innotech densitometer.

DNA Analysis

Southern analyses of genomic DNA was as described in Weeks et al., supra, except that 10 ug of DNA were loaded into each

slot of a 0.6% agarose gel. The probe consisted of the 1578 bp PvuII fragment from the coding region of Dx5 (bracketed in Figure 2B).

RESULTS

Forty-five independent callus pieces were selected from 4980 embryos bombarded in 10 different experiments; each yielded one to five plants. The T₀ primary regenerated plants and the lines derived from them are designated with a unique number following the bombardment number, eg., plant 126 regenerated from bombardment 7 is 7-126. Plants from twenty-six of these independent lines gave rise to progeny embryos that were able to germinate in the presence of 3 mg/L bialaphos when excised from immature seeds three to four weeks after anthesis.

The immature endosperm tissue corresponding to each excised embryo was extracted and screened by SDS-PAGE for the presence of the hybrid glutenin. This SDS-PAGE analysis showed that the migration of the hybrid HMW glutenin subunit is clearly distinguishable from the native HMW subunits under these gel conditions.

Although the lanes did not contain the same amounts of total protein, it was evident that there is variation in the expression levels of the hybrid subunit relative to those of the native HMW subunits in the same seed. Of the twenty-six resistant lines that were derived from co-bombardment experiments including pGlut10H5, sixteen (60%) lines showed expression of the hybrid subunit in the T₁ generation.

Expression stability

At each successive generation, 24 or 25 immature embryos 3-4 weeks after anthesis were tested for their ability to germinate on bialaphos. Plants were classified as homozygous if all 25 of their progeny embryos exhibited resistance. The levels of transgene expression were assessed in successive generations. In most cases, expression was stable. An apparent increase in expression between the T₁ and T₂ generations in some plants was explained by the increase in average gene dosage between the heterozygous and homozygous progeny of the T₀ plant and T₁ generation, respectively. Since the endosperm is triplicate, individual selfed progeny of the T₀

heterozygote could have 0, 1, 2 or 3 copies and the T₁ lane contains an average of 8 of these individuals.

In four of the 15 wheat lines analyzed, expression of the hybrid HMW glutenin has not remained at the same levels, but rather has declined as homozygous resistant lines were selected in the T₂ or the T₃ generations. The decline could be due either to hybrid HMW glutenin transgene inactivation or to loss of unlinked pGlu10H5 transgene(s) copies via segregation.

Cross-linking ability of the hybrid subunit

As noted above, the hybrid HMW glutenin contains a unique configuration of cysteines, γ-type at the N-terminus and α-type at the C-terminus. The hybrid HMW glutenin behaved exactly like the native subunits in terms of its solubility in SDS buffers with and without a reducing agent, when examined by SDS-PAGE. The native and hybrid HMW glutenins were partially extracted from both the Bobwhite and the transgenic flours by SDS alone and were present in lower molecular weight polymers. The majority of the hybrid glutenin, however, was insoluble in SDS buffer and could only be extracted when a reducing agent was added. Thus, the configuration of cysteines in the hybrid HMW glutenin does not preclude the formation of either intermolecular disulfide bonds or high molecular weight polymers.

Relative expression levels

Figure 3 shows SDS-PAGE analysis of protein extracts from eight different transgenic lines (Lanes 1-8) and Bobwhite (C). Each sample was prepared from flour of a mixture of eight mature seeds of a single homozygous T₂ or T₃ plant. Under these gel conditions, the native By9 and Dyl0 subunits are not well resolved and run as a closely spaced doublet of apparent molecular weight 94 kDa. Likewise the two isoforms of the Dx5 subunit are not well-resolved and migrate as a broad band. The Bx7 band is well-separated from other proteins and the darkest of the HMW subunits in intensity. In all the lines except the one shown in Lane 3, the hybrid HMW subunit is present in at least as high a quantity as the native Ax2* (largest) HMW subunit. Thus even a single copy of the hybrid transgene (Lane

1) can support significant expression levels, comparable to those of the native HMW subunits.

In one of the lines (shown in lane 2), synthesis of native HMW glutenin synthesis has declined relative to the other proteins. The extract shown there exhibits reduced synthesis of all the native HMW glutenin compared to the lower molecular weight seed proteins in the same lane.

In order to estimate the levels of the hybrid HMW glutenin relative to those of the native HMW glutenin and other seed proteins, the gel was scanned by densitometry. All the wheat prolamines stay on the gel. However, since proteins of molecular weights less than 33 kDa are run off the gel under the conditions used to separate the hybrid HMW glutenin from the nearby Dx5 subunit band, only relative quantities could be assessed. The results of these comparisons are shown in Table I along with the number of intact transgene copies estimated from restriction enzyme digests of genomic DNA (see below). The hybrid HMW glutenin band constitutes more than 20% of the total HMW glutenins in all the lines except that in lane 3. Thus the transgene has made a significant contribution to the HMW glutenin composition in all the transgenic lines.

In the case where endogenous HMW glutenin accumulation is reduced (Lane 2), the hybrid makes up 62% of the total complement of HMW glutenin. In most lines, the sum of the HMW glutenin accumulation is elevated 29 to 45% relative to the HMW glutenin content of Bobwhite (all sums are normalized to the remainder of the protein in their respective lanes). Thus this quantitative analysis suggests that addition of HMW-glutenin gene copies raises the levels of HMW glutenin accumulation in mature seeds.

DNA analysis

DNA was isolated from the leaves of transgenic plants of each line and assayed for the presence of transgenes by genomic Southern analysis, as described above. Hybridizing bands known to correspond to the transgenes were scanned and their densities compared. The results are shown in Table I. The number ranges from 1 in line 11-291 to about 6 in 14-23 (Table I). This range is much broader than that in total HMW glutenin levels or even hybrid HMW glutenin levels. However the most

highly expressing line had one of the highest copy numbers (Line 4) and the poor expression was in a single copy line (Line 3).

Table I. Transgenic HMW-glutenin gene activity in individual wheat lines.

Line	Transgene Expression	Total subunit as % of total	HMW Gene Copy No. Estimate
1	23%	135%	1
2	62%	70%	5-6
3	15%	145%	1
4	42%	145%	4-5
5	23%	140%	2
6	24%	129%	2
7	22%	135%	3-4
8	20%	143%	3-4

DISCUSSION

The experiments reported here show that levels of HMW glutenins can be increased by addition of gene copies. The behavior of the hybrid subunit is indistinguishable from that of the native subunits. The protein accumulates in the endosperm tissue over the course of seed development and is cross-linked by disulfide bonds into large polymers that cannot be extracted by SDS without addition of reducing agents.

Expression of the hybrid glutenin is under the control of the native Dyl0 5' and Dx5 3' flanking sequences. These sequences were sufficient to achieve high levels of the modified seed storage protein. Even a single gene copy supported expression levels comparable to those of the native HMW glutenin genes. Thus, these sequences serve as a source of effective regulatory sequences suitable for expression of other

proteins in transgenic wheat and potentially other cereal endosperm.

Example 2

This example describes the transformation of wheat using a recombinant construct encoding Dx5 and Dyl0.


The recombinant expression vectors were prepared as generally as described above except that the complete Dx5 or Dyl0 gene, including 5' and 3' regulatory sequences, were used. The constructs were then introduced into immature wheat embryos by particle bombardment as described above. Analysis of T₀ plants and their progeny is performed as described above.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

Applicant's or agent's file reference number A700/363	International application No. ...
--	-----------------------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>17</u> , line <u>31</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution Agricultural Research Culture Collection (NRRL)	
Address of depositary institution (including postal code and country) 1815 North University Street Peoria, Illinois 61604 US	
Date of deposit January 12, 1996	Accession Number NRRL B-21517
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
1 microorganism is disclosed in the specification. In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication or the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<input checked="" type="checkbox"/> For receiving Office use only This sheet was received with the international application	<input type="checkbox"/> For International Bureau use only This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer

WHAT IS CLAIMED IS:

1. A method of altering glutenin content in the endosperm of a wheat plant, the method comprising:

stably transforming a parental wheat plant with a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and a seed-specific promoter to produce a stably transformed wheat plant; and

selecting a progeny of said stably transformed wheat plant having a glutenin content in the endosperm of a mature seed at least about 30% different than the glutenin content in the endosperm of a mature seed from the parental wheat plant.

2. The method of claim 1, wherein the nucleic acid is stably transformed into the first wheat plant using particle bombardment.

3. The method of claim 1, wherein the nucleic acid encodes a chimeric glutenin polypeptide.

4. The method of claim 1, wherein the chimeric polypeptide comprises sequences from an x-type glutenin polypeptide and a y-type glutenin polypeptide.

5. The method of claim 1, wherein the x-type polypeptide is encoded by the Glu-D1-1b gene.

6. The method of claim 1, wherein the y-type polypeptide is encoded by the Glu-D1-2b gene.

7. The method of claim 1, wherein the promoter is from the Glu-D1-2b gene.

8. The method of claim 1, wherein the parental wheat plant is Bobwhite.

9. A wheat plant comprising a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and a seed-specific promoter, the plant having a

glutenin content in the endosperm of a mature seed at least about 30% different than the glutenin content in the endosperm of a mature seed from a parental wheat plant.

10. The plant of claim 9, wherein the nucleic acid encodes a chimeric glutenin polypeptide.

11. The plant of claim 10, wherein the chimeric polypeptide comprises sequences from an x-type polypeptide and a y-type glutenin polypeptide.

12. The plant of claim 11, wherein the x-type polypeptide is encoded by the Glu-D1-1b gene.

13. The plant of claim 11, wherein the y-type polypeptide is encoded by the Glu-D1-2b gene.

14. The plant of claim 9, wherein the promoter is from the Glu-D1-2b gene.

15. The plant of claim 9, wherein the high molecular weight (HMW) glutenin accounts for at least about 15% of the total protein in the endosperm of the mature seed.

16. The plant of claim 9, wherein the parental wheat plant is Bobwhite.

17. A seed derived from the plant of claim 9.


18. A recombinant construct comprising a wheat glutenin gene promoter of about 400 to about 2800 nucleotides operably linked to a heterologous polynucleotide sequence encoding a chimeric glutenin polypeptide, wherein the chimeric glutenin polypeptide is encoded by polynucleotide sequences from the Glu-D1-1b gene and the Glu-D1-2b gene.

19. The recombinant construct of claim 18, wherein the wheat glutenin gene is Glu-D1-2b.

20. A plant comprising the recombinant construct of claim
18.

21. Wheat flour derived from the plant of claim 1.

Y-type HMW-glutenin



1/3

X-type HMW-glutenin






FIG. 1

2/3

**FIG. 2A****FIG. 2B**

3/3




FIG. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/00510

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/200, 205, 250, DIG 58; 435/172.3, 240.4, 320.1; 536/23.6, 24.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CABA, MEDLINE, CAPLUS, GENBANK, BIOSIS

search terms: wheat, glutenin, transformation, Glu-D1-2b, Glu-D1-1b

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SHANI et al. Role of amino- and carboxy- terminal regions in the folding and oligomerization of wheat high molecular weight glutenin subunits. Plant Physiol. 1992, Vol. 98, pages 433-441, see entire document.	1-21
Y	WEEKS et al. Rapid production of multiple independent lines of fertile transgenic wheat (<i>Triticum aestivum</i>). Plant Physiol. 1993, Vol. 102, pages 1077-1084, especially pages 1078-1079.	1-21
Y	ANDERSON et al. Nucleotide sequences of the two high-molecular-weight glutenin genes from the D-genome of a hexaploid bread wheat, <i>Triticum aestivum</i> L. cv Cheyenne. Nucleic Acids Research. 1989, Vol. 17, No. 1, pages 461-462, see entire document.	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

20 MARCH 1997

Date of mailing of the international search report

08 APR 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimil No. (703) 305-3230

Authorized officer

THOMAS HAAS

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/00510

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Categ ry*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>GALILI, G. Heterologous expression of a wheat high molecular weight glutenin gene in <i>Escherichia coli</i>. Proc. Natl. Acad. Sci. USA. October 1989, Vol. 86, pages 7756-7760, see entire document.</p>	1-21

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/00510

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12N 15/00, 15/05, 15/09, 15/11, 15/29, 15/64, 15/82; A01H 1/00, 3/00, 4/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :


800/200, 205, 250, DIG 58; 435/172.3, 240.4, 320.1; 536/23.6, 24.1



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00, 15/05, 15/09, 15/11, 15/29, 15/64, 15/82, A01H 1/00, 3/00, 4/00		A1	(11) International Publication Number: WO 97/25419
(21) International Application Number: PCT/US97/00510		(43) International Publication Date: 17 July 1997 (17.07.97)	
(22) International Filing Date: 10 January 1997 (10.01.97)		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
(30) Priority Data: 08/586,331 12 January 1996 (12.01.96) US			
(71) Applicant: THE UNITED STATES OF AMERICA, represented by TH [US/US]; E SECRETARY OF AGRICULTURE, Washington, DC 20250 (US).			
(72) Inventors: BLECHL, Ann, E.; 1005 Buchanan Street, Albany, CA 94706 (US). ANDERSON, Olin, D.; 208 Poshard Street, Pleasant Hill, CA 94523 (US).			
(74) Agents: PENDORF, Stephan, A. et al.; Suite 1000, 600 N. Westshore Boulevard, Tampa, FL 33609 (US).			

(54) Title: GLUTENIN GENES AND THEIR USES



(57) Abstract

The present invention is directed to methods of improving flour by altering glutenin content in seeds of plants, particularly wheat. The invention relates to methods of introducing a recombinant construct comprising a glutenin gene into a parental plant.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

GLUTENIN GENES AND THEIR USES

Field of the Invention

The present invention relates generally to plant molecular biology. In particular, it relates to nucleic acids and methods for improving glutenin content of plants.

BACKGROUND OF THE INVENTION

The glutenins, which include both high molecular weight (HMW) glutenin subunits and low molecular weight (LMW) glutenin subunits, comprise an economically important class of wheat seed storage proteins. The apparent molecular weights of the individual HMW glutenin polypeptides or subunits range from 90 to 200 kDa. These subunits crosslink by disulfide bonds among themselves and with LMW glutenin polypeptides to form polymers exceeding one million daltons in molecular weight. HMW glutenins constitute 8-10%, while LMW glutenins constitute 15-20% of the total endosperm protein. Both HMW and LMW glutenin proteins play important functional roles in determining the end-uses of wheat flour.

In wheat, HMW glutenins are encoded at the Glu-1 loci on the long arms of the group 1 chromosomes. Each locus consists of two separate genes, encoding an x-type and a y-type subunit, respectively. These pairs have never been confirmed to be separated by recombination. This has made determination of their separate contributions to bread dough properties difficult to assess by genetic correlation studies. For a review of the genetics and biochemistry of glutenin polypeptides, see, Shewry et al., J. Cereal Sci. 15:105-120 (1992).

Both the quantity and identity of specific HMW glutenin alleles contribute to the differences in bread-making quality of various cultivars. For instance, deletion of glutenin genes results in a decrease in the overall levels of HMW glutenins, which results in decreases in bread-making quality (see, e.g., Lawrence et al. J. Cereal Sci 7:109-112 (1988)).

The effects of overproducing HMW glutenin on protein accumulation and baking quality has not been assessed because such lines of wheat have not been found among natural

populations. In addition, direct alteration of the glutenin subunits that form the polymers is not possible using standard breeding methods. Thus, the art lacks reproducible and efficient methods of producing lines with altered glutenin contents. The present invention addresses these and other needs.

SUMMARY OF THE INVENTION

The present invention provides methods of increasing glutenin in the endosperm of wheat plants. The methods and plants of the invention therefore are useful in providing flour and dough having improved end-use properties. The methods comprise introducing into a parental wheat plant a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and selecting progeny wheat plant having increased glutenin content in the endosperm of mature seed. The glutenin content of the progeny is preferably at least about 15% greater than the glutenin of the parental wheat plant.

Any method of introducing the expression cassette into the parental plant can be used. Particle bombardment is a convenient method for producing transgenic plants. Once the expression cassette is stably integrated into the genome, standard sexual crosses can be used to introduce the expression cassette into desired lines.

In some embodiments, the nucleic acid introduced into the plant encodes a chimeric glutenin polypeptide. For instance, the chimeric polypeptide may comprise sequences from an x-type glutenin polypeptide and a y-type glutenin polypeptide. Exemplary genes for this purpose include the Glu-D1-1b gene and the Glu-D1-2b gene.

The expression cassette may further comprise a seed-specific promoter to direct expression of the introduced nucleic acid to the endosperm. A convenient promoter for this purpose is the promoter from the Glu-D1-2b gene.

Any wheat cultivar can be used as the parental line in the present invention. An exemplary cultivar is Bobwhite.

The invention also provides wheat plants comprising a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide. The plants of the invention

have a glutenin content in the endosperm of a mature seed at least about 15% greater than the glutenin content in the endosperm of a mature seed from a parental wheat plant. The percentage of total endosperm protein which is glutenin will usually depend upon the glutenin content of the parental line. Usually, HMW glutenins account for at least about 15% of the total protein in the endosperm of the mature seed from plants of the invention.

The invention further provides recombinant constructs comprising a wheat glutenin gene promoter of about 400 to about 2800 nucleotides operably linked to a heterologous polynucleotide sequence. These constructs are particularly useful in directing expression of the heterologous sequence to seeds of transgenic plants. An exemplary wheat glutenin gene from which the promoters of the invention can be derived is Glu-D1-2b.

Definitions

The term "plant" includes whole plants, plant organs (e.g., leaves, stems, roots, flowers, etc.), seeds and plant cells and progeny of same. The class of plants which can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

A "heterologous sequence" is one that originates from a foreign species, or, if from the same species, is substantially modified from its original form. For example, a promoter operably linked to a heterologous structural gene is from a species different from that from which the structural gene was derived, or, if from the same species, one or both are substantially modified from their original form.

A "glutenin polypeptide" is a gene product of a glutenin gene or glutenin polynucleotide sequence. A glutenin polypeptide can be either a LMW glutenin or a HMW glutenin. A glutenin polypeptide contains cysteine residues by which disulfide bonds are formed with other glutenin polypeptides to form polymers. The composition and size of the repeat region is also important to polymer formation.

A "chimeric glutenin polypeptide" is glutenin gene product that comprises a modified amino acid sequence. Modifications, as explained in detail below, can be in the form of substitutions, deletions, or additions of single amino acids or groups of amino acids. Thus, chimeric glutenin polypeptides can be hybrid glutenin polypeptides comprising sequences from two or more different subunits or may be polypeptides in which single amino acid modifications are made.

In the case where an inserted polynucleotide sequence is transcribed and translated to produce a functional glutenin polypeptide, one of skill will recognize that because of codon degeneracy, a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the terms "glutenin gene" or "glutenin polynucleotide sequence". In addition, the terms specifically include those full length sequences substantially identical (determined as described below) with a glutenin gene sequence and that encode proteins that retain the function of the glutenin polypeptide. Thus, in the case of wheat glutenin genes disclosed here, the term includes variant polynucleotide sequences which have substantial identity with the sequences disclosed here and which encode glutenin polypeptides capable of crosslinking by disulfide bonds with other glutenin polypeptides to form glutenin polymers.

Two polynucleotides or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the complementary sequence is identical to all or a portion of a reference polynucleotide sequence.

Sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a segment or "comparison window" to identify and compare local regions of sequence similarity. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman Adv. Appl. Math. 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch J. Mol. Biol. 48:443 (1970), by the search for similarity method of Pearson and Lipman Proc. Natl.

Acad. Sci. (U.S.A.) 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection. "Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 60% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using the programs described above using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 40%, preferably at least 60%, more preferably at least 90%, and most preferably at least 95%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids

having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C to about 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. However, nucleic acids which do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a diagram showing the domains of γ -type and α -type HMW glutenins. The C- and N-terminal domains are shaded, the locations of cysteine residues are designated by S.

Figure 2A is a diagram showing the structure of a hybrid glutenin polypeptide of the invention.

Figure 2B is a diagram of the recombinant plasmid of the invention which encodes the polypeptide shown in Figure 2A.

Figure 3 is a photograph of a gel stained with Coomassie blue showing the relative mobility of a hybrid HMW glutenin polypeptide of the invention (marked by the arrow).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

This invention relates to plant glutenin genes, in particular wheat glutenin genes. Nucleic acid sequences from glutenin genes can be used to modify glutenin content in transgenic plants, in particular wheat plants. In addition,

the invention provides new seed-specific promoters useful for directing expression of desired genes in a number of plants.

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning - A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

Glutenin polypeptides and genes

The HMW glutenins are composed primarily of a central domain of 45-90 repeating two or three simple motifs, comprising chiefly glutamine and proline. This unusual primary structure results in a rod-like secondary structure as assessed by circular dichroism and SEM. Cysteines in the C- and N-terminal domains of the molecules are known to be critical for the formation of intermolecular disulfide bonds (Figure 1). The quaternary interactions of these proteins are of interest because the disulfide crosslinks among HMW and LMW subunits is thought to be integral to the elastic properties of dough.

Any isolated glutenin gene can be used in the present invention. The particular polynucleotide sequence used is not a critical feature of the invention, so long as the desired alteration in glutenin content is achieved. As noted above, HMW glutenins (both x-types and y-types) are encoded at the Glu-1 locus. In hexaploid wheat the group 1 chromosomes are designated 1A, 1B and 1D. Studies of the frequency of alleles at each of the loci indicate the presence of at least three alleles at Glu-A1, 11 alleles at Glu-B1, and six alleles at Glu-D1. The gene products of each allele are designated by noting the chromosome, followed by its classification as x or y, and a number (see, MacRitchie, Advances in Food and Nutrition Research 36:1-87 (1992)).

Wheat HMW glutenin genes have been cloned and described in the literature. For instance, a complete set of HMW glutenin

genes have been isolated and sequenced from the hard red winter wheat Cheyenne (Anderson et al. Wheat Genetics Symposium Proc. pp 699-704 (Bath Press, Cambridge, UK, 1988), including two, Dx5 and Dyl0, with the highest correlation with good flour quality (Anderson et al. Nuc. Acids. Res. 17:461-462 (1989)). Other particularly useful HMW glutenins include Ax and Bx alleles.

The isolation of other glutenin genes may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed in the prior art can be used to isolate the desired gene from a cDNA or genomic DNA library. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a cDNA library, mRNA is isolated from endosperm and a cDNA library which contains the glutenin gene transcript is prepared from the mRNA.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology to amplify the sequences of the glutenin and related genes directly from genomic DNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds., Academic Press, San Diego (1990)).

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers et al., Cold Spring Harbor Symp. Quant. Biol. 47:411-418 (1982), and Adams et al., J. Am. Chem. Soc. 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding

the complementary strand using DNA polymerase with an appropriate primer sequence.

Isolated sequences prepared as described herein can then be used to modify glutenin gene expression and therefore glutenin content in plants. One of skill will recognize that the nucleic acid encoding a functional glutenin protein need not have a sequence identical to the exemplified genes disclosed here. Thus, genes encoding chimeric glutenin polypeptides can be used in the present invention.

As noted above, glutenin polypeptides, like other proteins, have different domains which perform different functions. Thus, the glutenin gene sequences need not be full length, so long as the desired functional domains of the protein is expressed. Chimeric glutenin polypeptides can be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art. For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like. In particular, cysteine residues may be added, deleted or moved within the polypeptide to achieve a modified glutenin polypeptide with desired properties. Chimeric polypeptides may also be produced by fusing coding sequences from two or more glutenin genes. All of these modifications can be used in a number of combinations to produce the final modified protein chain.

Preparation of recombinant constructs

To use isolated glutenin sequences in modifying glutenin content in plants, recombinant DNA vectors suitable for transformation of plant cells are prepared. A DNA sequence coding for the desired glutenin polypeptide, for example a cDNA or a genomic sequence encoding a full length protein, is conveniently used to construct a recombinant expression cassette which can be introduced into the desired plant. An expression cassette will typically comprise the glutenin polynucleotide sequence operably linked to a promoter sequence and other transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the glutenin gene in the intended tissues (e.g., endosperm) of the transformed plant.

For example, a constitutive plant promoter fragment may be employed which will direct expression of the glutenin in all tissues of a plant. Such promoters are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may be under environmental control. Such promoters are referred to here as "inducible" promoters. Examples of environmental conditions that may effect transcription by inducible promoters include pathogen attack, anaerobic conditions, or the presence of light.

Typically, the promoters used in the constructs of the invention will be "tissue-specific" and are under developmental control such that the desired gene is expressed only in certain tissues, such as leaves, roots, fruit, seeds, or flowers. Promoters that direct expression in seeds, particularly the endosperm are particularly preferred. Examples of such promoters include the promoter from genes encoding seed storage proteins, such as napin, cruciferin, phaseolin, and the like (see, U.S. Patent No. 5,420,034). Other promoters suitable for expressing glutenin genes in cereals include promoters from genes encoding gliadins, cereal prolamines (e.g., zein, hordein, secalin, and avenin) and starch biosynthetic enzymes.

The endogenous promoters from glutenin genes are particularly useful for directing expression of glutenin genes to the seed, particularly the endosperm. These seed-specific promoters can also be used to direct expression of heterologous structural genes. Thus, the promoters can be used in recombinant expression cassettes to drive expression of any gene whose expression in seeds is desirable. Examples include genes encoding proteins useful in increasing the nutritional value of seeds (e.g., genes encoding proteins involved in lipid, protein, and carbohydrate or starch biosynthesis). Other genes include those encoding pharmaceutically useful

compounds, and genes encoding plant resistance products to combat fungal or other infections of the seed.

The glutelin promoters can also be used to initiate transcription of mRNA molecules to inhibit expression of an endogenous endosperm gene. Exemplary genes whose expression could be inhibited include genes encoding glutenins, enzymes involved in carbohydrate or lipid synthesis, secalins, and the like. Means for inhibiting gene expression in plants using recombinant DNA techniques are well known. For instance, antisense technology can be conveniently used. see, e.g., Sheehy et al., Proc. Nat. Acad. Sci. USA, 85:8805-8809 (1988), and Hiatt et al., U.S. Patent No. 4,801,340. Catalytic RNA molecules or ribozymes can also be used to inhibit expression of endosperm-specific genes. The design and use of target RNA-specific ribozymes is described in Haseloff et al. Nature, 334:585-591 (1988). Introduction of nucleic acid configured in the sense orientation has also been shown to be an effective means by which to block the transcription of target genes. For an example of the use of sense suppression to modulate expression of endogenous genes see, Napoli et al., The Plant Cell 2:279-289 (1990), and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

The glutenin promoters of the invention are typically at least about 400 base pairs in length, and often at least about 800 or about 1000 base pairs. The length of the promoters is typically less than about 3500 base pairs, usually less than about 2800 base pairs and often less than about 2000 base pairs in length. The length of the promoters is counted upstream from the translation start codon of the native gene. One of skill will recognize that use of the "about" to refer to lengths of nucleic acid fragments is meant to include fragments of various lengths that do not vary significantly from the lengths recited here and still maintain the functions of the claimed promoters (i.e., seed-specific gene expression).

To identify glutenin promoters, the 5' portions of a genomic glutenin gene clone is analyzed for sequences characteristic of promoter sequences. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In plants, further upstream from the

TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing et al., in Genetic Engineering in Plants, pp. 221-227 (Kosage, Meredith and Hollaender, eds. 1983).

In preparing expression vectors of the invention, sequences other than the promoter and the structural gene of interest are also preferably used. If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the glutenin coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences from a glutenin gene will typically comprise a marker gene which confers a selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron, or phosphinothricin (the active ingredient in bialaphos and Basta).

Preparation of transgenic plants

The DNA constructs described above may be introduced into the genome of the desired plant host by a variety of conventional techniques. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising et al., Ann. Rev. Genet. 22:421-477 (1988).

The DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as biolistic methods, electroporation, PEG poration, and microinjection of plant cell protoplasts or embryogenic callus. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced using an *Agrobacterium tumefaciens* or *A. rhizogenes* vector.

Particle bombardment techniques are described in Klein et al., Nature 327:70-73 (1987). A particularly preferred method of transforming wheat and other cereals is the bombardment of calli derived from immature embryos as described by Weeks et al. Plant Physiol. 102:1077-1084 (1993).

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984). Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA* 82:5824 (1985).

Agrobacterium tumefaciens-mediated transformation techniques are also well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983). Although *Agrobacterium* is useful primarily in dicots, certain monocots can be transformed by *Agrobacterium*. For instance, *Agrobacterium* transformation of rice is described by Hiei et al. *Plant J.* 6:271-282 (1994).

The present invention is particularly useful in wheat and other cereals. A number of methods of transforming cereals have been described in the literature. For instance, transformation of rice is described by Toriyama et al. *Bio/Technology* 6:1072-1074 (1988), Zhang et al. *Theor. Appl. Gen.* 76:835-840 (1988), and Shimamoto et al. *Nature* 338:274-276 (1989). Transgenic maize regenerants have been described by Fromm et al., *Bio/Technology* 8:833-839 (1990) and Gordon-Kamm et al., *Plant Cell* 2:603-618 (1990)). Similarly, oats (Sommer et al., *Bio/Technology* 10:1589-1594 (1992)), wheat (Vasil et al., *Bio/Technology* 10:667-674 (1992)); Weeks et al., *Plant Physiol.* 102:1077-1084 (1993)), sorghum (Casas et al., *Proc. Natl. Acad. Sci. USA* 90:11212-11216 (1993)), rice (Li et al., *Plant Cell Rep.* 12:250-255 (1993)), barley (Yuechun and Lemaux, *Plant Physiol.* 104:37-48 (1994)), and rye (Castillo et al., *Bio/Technology* 12:1366-1371 (1994)) have been transformed via bombardment.

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the glutenin polynucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation*

and Culture, Handbook of Plant Cell Culture, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, Regeneration of Plants, Plant Protoplasts, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. Ann. Rev. of Plant Phys. 38:467-486 (1987).

The methods of the present invention are particularly useful for incorporating the glutenin polynucleotides into transformed plants in ways and under circumstances which are not found naturally. In particular, the glutenin polypeptides may be expressed at times or in quantities which are not characteristic of natural plants.

One of skill will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

The invention has use over a broad range of types of plants. The glutenin genes are preferably expressed in cereal species commonly used for production of flour, e.g., wheat, rye, oats, and the like.

The seed-specific promoters from glutenin genes can be used in essentially any plant species. For instance, the promoters or genes described here can be used in species from the genera Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Cucumis, Cucurbita, Daucus, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Juglans, Lactuca, Linum, Lolium, Lycopersicon, Malus, Manihot, Majorana, Medicago, Nicotiana, Oryza, Panium, Pannisetum, Persea, Phaseolus, Pisum, Pyrus, Prunus, Raphanus, Secale, Senecio, Sinapis, Solanum, Sorghum, Trigonella, Triticum, Vitis, Vigna, and, Zea. Plants that are particularly useful in the invention include Avena, Brassica, Glycine, Hordeum, Oryza, Phaseolus, Pisum, Secale, Sorghum, Triticum, Vigna, and Zea.

As noted above, the invention is particularly useful for improvement of wheat cultivars. Any wheat cultivar may be improved by the methods of the invention. Exemplary lines

include hexaploid lines such as Anza, Shasta, Yecoro Rojo, Siouxland, Freedom, Tam107, and Tam200.

The effect of the modification of glutenin gene expression can be measured by detection of increases or decreases in desired glutenin protein levels using, for instance, gel electrophoresis, as described below. Quantification of HMW glutenin content can be carried out by SDS/PAGE densitometry, as described below.

Other methods for quantifying glutenin content include sonication of flour dispersions in SDS buffer to solubilize glutenin and other relatively insoluble proteins (see, Singh and MacRitchie in Wheat End-Use Properties: Wheat and Flour Characterization for Specific End-Uses H. Salovaara, ed., pp 321-326 (University of Helsinki, Helsinki, 1989)). This method is useful in solubilizing at least 95% of total flour protein.

Quantification of the glutenin fraction can then be performed using size exclusion high performance liquid chromatography as described by Batey et al., Cereal Chem. 68:207-209 (1991). Other suitable methods include reverse phase HPLC and capillary electrophoresis.

The plants of the invention have increased in glutenin content in the endosperm of mature seeds as compared to the parental lines from which they are derived. The increase is usually at least about 15% as compared to the parental line. More usually, the increase is at least about 20%, preferably at least about 35%, and in more preferred embodiments at least about 45%. Any of the methods for quantifying glutenins in the endosperm described above can be used to determine the percent increase in glutenin content. A wheat seed typically reaches maturity about 6 to about 10 weeks after anthesis.

Another measure of increased glutenin content in transgenic plants of the invention is glutenin content as a percent of total protein of a mature seed or flour derived from the mature seed. Different wheat cultivars have different glutenin contents as measured using standard methods. Thus, the final glutenin content of the plants of the invention will depend upon the parental line. In the case where HMW glutenin genes are used, the plants of the invention will typically have HMW glutenin contents of at least about 15%, usually at least

about 18%, and preferably at least about 20% as measured using standard methods such as those described in detail below.

Analysis of the rheological properties of flour derived from the transgenic plants of the invention can be carried out according to standard physical dough-testing instruments widely used to measure flour and dough quality (see, e.g., MacRitchie, *Advances in Food and Nutrition Research* 36:1-87 (1992)). Such methods include, for instance, use of extensographs to measure tensile strength. Two of the main parameters measured are maximum resistance (R_{max}) and extensibility (Ext). Other methods include mixographs and bake-test loaf volume (MacRitchie and *Gras Cereal Chem.* 50:292-302 (1973) as well as SDS-sedimentation tests, amelographs, and cookie spread methods.

The following Examples are offered by way of illustration, not limitation.

Example 1

This example demonstrates increase in glutenin content in transgenic wheat plants comprising recombinant constructs of the invention.

MATERIALS AND METHODS

Construction of the hybrid HMW glutenin expression plasmid

Two HMW glutenin subunits, Dx5 and Dy10, encoded by the Glu-D1-1b and Glu-D1-2b genes, respectively, have been most often associated with superior dough strength by genetic correlation studies. A bacterial expression vector comprising a fusion of coding regions of the Dy10 and Dx5 genes is described by Shani et al., *Plant Physiol.* 98:433-441 (1992). The hybrid HMW glutenin polypeptide encoded by this gene is diagrammed in Figure 2A.

In this example, the Dy10:Dx5 hybrid coding region was reunited with 5' and 3' flanking sequences from the native Dy10 and Dx5 genes. The construction of a fusion at the HindIII site located at base pairs 429-435 and 385-391 after the A of the start codons of the Glu-D1-2b (Genbank Accession N. X12929) and Glu-D1-1b (Genbank Accession No. X12928) genes (Anderson et al. *Nuc. Acids Res.* 17:461-462 (1989)) and its

insertion into the bacterial expression vector pet3A to form pet-3a-10/5 is described by Shani et al., supra.

Thus, the native Dyl0 gene provides the promoter, transcription start site, the 5' transcribed untranslated region and the first 145 codons including the 21 amino acid signal peptide. The native Dx5 gene provides its codons 130 to 848 followed by two translation stop codons, the transcription termination and poly(A) addition signals. Because amino acids 138 through 147 are shared by the Dyl0 and Dx5 glutenin subunits, the central region of repeating amino acid motifs of the hybrid glutenin polypeptide (thickest boxes in Figure 2A) is identical to that of Dx5. The configuration of the cysteines (S) and thus of potential disulfide bonds is chimeric in nature: the N-terminal cysteines are of the y-type and the single C-terminal cysteine is typical of x-type subunits.

In order to reconstitute the promoter region, transcription start and termination sites, and any other regulatory signals needed for expression in wheat, three fragments were isolated and combined in a single ligation reaction with the Bluescript KS⁻ (Stratagene) vector cut with EcoRI and BamHI: 1) the 715 bp StuI/DrdI fragment that contains the junction region in pET-3a-10/5; 2) the 2800 bp fragment from a clone of the native Dyl0 gene beginning at the EcoRI site 5' to the gene and ending at the StuI site 74 bp after the A in the start codon; and 3) the 3800 bp fragment from a clone of the native Dx5 gene that starts at the DrdI site 744 bp after the A in the start codon and ends at a BamHI site in the vector just outside the EcoRI site in the 3' flanking region. A plasmid with the correct structure was identified by restriction analysis and named pGlu10H5 (Figure 2B). This plasmid was deposited under the terms of the Budapest Treaty in the Agricultural Research Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois, USA, on January 12, 1996 and has been assigned Accession No. NRRL B-21517.

The plasmid UBI:BAR which comprises the maize ubiquitin promoter operably linked to the bar gene encoding the enzyme PAT, which inactivates phosphinothricin was prepared as described Corn jo et al. Plant Mol. Biol. 23:567-581 (1993).

Both plasmids were prepared by the alkaline lysis method using a Qiagen kit for the final purification. The DNAs were stored in TE buffer at a concentration of 1 mg/ml.

Wheat Transformation

12.5 ug of each DNA (an approximately 2:1 molar ratio of UBI:BAR to pGlul0H5) was coated onto gold particles for bombardment into immature wheat embryos of cultivar Bobwhite essentially as described by Weeks et al. Plant Physiol. 102:1077-1084 (1993) except that the embryos were incubated on callus induction media containing mannitol for 4 hours before and 20 hours after bombardment. Transformants were selected on the basis of their resistance to 1mg/L bialaphos at the callus and green shoot stages of regeneration and to 3 mg/L bialaphos at the rooting stage of regeneration as described by Weeks et al., supra.

Protein Analysis

Protein extracts from immature wheat endosperm were prepared by grinding the tissue or cells in SDS-PAGE sample loading buffer [66 mM Tris pH 6.8, 3% (w/v) SDS, 0.05% (w/v) Bromphenol blue, 2% (v/v) β -mercaptoethanol]. Protein extracts from mature dry wheat seeds were prepared by dissolving flour in sample buffer. SDS-PAGE was according to standard procedures. The separation gels for detection and quantitation of the hybrid subunit were 10% (w/v) acrylamide 0.05% (w/v) bis-acrylamide. The gels were run at 40 mamps constant current until 20 minutes after the bromphenol blue dye front had run off the gel (about 1000 Volt-hours).

The gels were stained by the Neuhoff method with a colloidal suspension of Coomassie blue. Protein standards for molecular weight calibration were purchased from Pharmacia and consisted of Phosphorylase b (94 kDa) Bovine Serum Albumin (67 kDa), Ovalbumin (43 kDa), Carbonic Anhydrase (30 kDa), Soybean Trypsin Inhibitor (20.1 kDa) and α -Lactalbumin (14.4 kDa). Stained gels were scanned using an Alpha Innotech densitometer.

DNA Analysis

Southern analyses of genomic DNA was as described in Weeks et al., supra, except that 10 ug of DNA were loaded into each

slot of a 0.6% agarose gel. The probe consisted of the 1578 bp PvuII fragment from the coding region of Dx5 (bracketed in Figure 2B).

RESULTS

Forty-five independent callus pieces were selected from 4980 embryos bombarded in 10 different experiments; each yielded one to five plants. The T₀ primary regenerated plants and the lines derived from them are designated with a unique number following the bombardment number, eg., plant 126 regenerated from bombardment 7 is 7-126. Plants from twenty-six of these independent lines gave rise to progeny embryos that were able to germinate in the presence of 3 mg/L bialaphos when excised from immature seeds three to four weeks after anthesis.

The immature endosperm tissue corresponding to each excised embryo was extracted and screened by SDS-PAGE for the presence of the hybrid glutenin. This SDS-PAGE analysis showed that the migration of the hybrid HMW glutenin subunit is clearly distinguishable from the native HMW subunits under these gel conditions.

Although the lanes did not contain the same amounts of total protein, it was evident that there is variation in the expression levels of the hybrid subunit relative to those of the native HMW subunits in the same seed. Of the twenty-six resistant lines that were derived from co-bombardment experiments including pGlut10H5, sixteen (60%) lines showed expression of the hybrid subunit in the T₁ generation.

Expression stability

At each successive generation, 24 or 25 immature embryos 3-4 weeks after anthesis were tested for their ability to germinate on bialaphos. Plants were classified as homozygous if all 25 of their progeny embryos exhibited resistance. The levels of transgene expression were assessed in successive generations. In most cases, expression was stable. An apparent increase in expression between the T₁ and T₂ generations in some plants was explained by the increase in average gene dosage between the heterozygous and homozygous progeny of the T₀ plant and T₁ generation, respectively. Since the endosperm is triplicate, individual selfed progeny of the T₀

heterozygote could have 0, 1, 2 or 3 copies and the T1 lane contains an average of 8 of these individuals.

In four of the 15 wheat lines analyzed, expression of the hybrid HMW glutenin has not remained at the same levels, but rather has declined as homozygous resistant lines were selected in the T₂ or the T₃ generations. The decline could be due either to hybrid HMW glutenin transgene inactivation or to loss of unlinked pGlut0H5 transgene(s) copies via segregation.

Cross-linking ability of the hybrid subunit

As noted above, the hybrid HMW glutenin contains a unique configuration of cysteines, y-type at the N-terminus and x-type at the C-terminus. The hybrid HMW glutenin behaved exactly like the native subunits in terms of its solubility in SDS buffers with and without a reducing agent, when examined by SDS-PAGE. The native and hybrid HMW glutenins were partially extracted from both the Bobwhite and the transgenic flours by SDS alone and were present in lower molecular weight polymers. The majority of the hybrid glutenin, however, was insoluble in SDS buffer and could only be extracted when a reducing agent was added. Thus, the configuration of cysteines in the hybrid HMW glutenin does not preclude the formation of either intermolecular disulfide bonds or high molecular weight polymers.

Relative expression levels

Figure 3 shows SDS-PAGE analysis of protein extracts from eight different transgenic lines (Lanes 1-8) and Bobwhite (C). Each sample was prepared from flour of a mixture of eight mature seeds of a single homozygous T₂ or T₃ plant. Under these gel conditions, the native By9 and Dyl0 subunits are not well resolved and run as a closely spaced doublet of apparent molecular weight 94 kDa. Likewise the two isoforms of the Dx5 subunit are not well-resolved and migrate as a broad band. The Bx7 band is well-separated from other proteins and the darkest of the HMW subunits in intensity. In all the lines except the one shown in Lane 3, the hybrid HMW subunit is present in at least as high a quantity as the native Ax2* (largest) HMW subunit. Thus even a single copy of the hybrid transgene (Lane

1) can support significant expression levels, comparable to those of the native HMW subunits.

In one of the lines (shown in lane 2), synthesis of native HMW glutenin synthesis has declined relative to the other proteins. The extract shown there exhibits reduced synthesis of all the native HMW glutenin compared to the lower molecular weight seed proteins in the same lane.

In order to estimate the levels of the hybrid HMW glutenin relative to those of the native HMW glutenin and other seed proteins, the gel was scanned by densitometry. All the wheat prolamines stay on the gel. However, since proteins of molecular weights less than 33 kDa are run off the gel under the conditions used to separate the hybrid HMW glutenin from the nearby Dx5 subunit band, only relative quantities could be assessed. The results of these comparisons are shown in Table I along with the number of intact transgene copies estimated from restriction enzyme digests of genomic DNA (see below). The hybrid HMW glutenin band constitutes more than 20% of the total HMW glutenins in all the lines except that in lane 3. Thus the transgene has made a significant contribution to the HMW glutenin composition in all the transgenic lines.

In the case where endogenous HMW glutenin accumulation is reduced (Lane 2), the hybrid makes up 62% of the total complement of HMW glutenin. In most lines, the sum of the HMW glutenin accumulation is elevated 29 to 45% relative to the HMW glutenin content of Bobwhite (all sums are normalized to the remainder of the protein in their respective lanes). Thus this quantitative analysis suggests that addition of HMW-glutenin gene copies raises the levels of HMW glutenin accumulation in mature seeds.

DNA analysis

DNA was isolated from the leaves of transgenic plants of each line and assayed for the presence of transgenes by genomic Southern analysis, as described above. Hybridizing bands known to correspond to the transgenes were scanned and their densities compared. The results are shown in Table I. The number ranges from 1 in line 11-291 to about 6 in 14-23 (Table I). This range is much broader than that in total HMW glutenin levels or even hybrid HMW glutenin levels. However the most

highly expressing line had one of the highest copy numbers (Line 4) and the poor st expression was in a single copy line (Line 3).

Table I. Transgenic HMW-glutenin gene activity in individual wheat lines.

Line	Transgene Expression	Total subunit as % of total	HMW Gene Copy No. Estimate
1	23%	135%	1
2	62%	70%	5-6
3	15%	145%	1
4	42%	145%	4-5
5	23%	140%	2
6	24%	129%	2
7	22%	135%	3-4
8	20%	143%	3-4

DISCUSSION

The experiments reported here show that levels of HMW glutenins can be increased by addition of gene copies. The behavior of the hybrid subunit is indistinguishable from that of the native subunits. The protein accumulates in the endosperm tissue over the course of seed development and is cross-linked by disulfide bonds into large polymers that cannot be extracted by SDS without addition of reducing agents.

Expression of the hybrid glutenin is under the control of the native Dyl0 5' and Dx5 3' flanking sequences. These sequences were sufficient to achieve high levels of the modified seed storage protein. Even a single gene copy supported expression levels comparable to those of the native HMW glutenin genes. Thus, these sequences serve as a source of effective regulatory sequences suitable for expression of other

proteins in transgenic wheat and potentially other cereal endosperm.

Exempl 2

This example describes the transformation of wheat using a recombinant construct encoding Dx5 and Dy10.

The recombinant expression vectors were prepared as generally as described above except that the complete Dx5 or Dy10 gene, including 5' and 3' regulatory sequences, were used. The constructs were then introduced into immature wheat embryos by particle bombardment as described above. Analysis of T₀ plants and their progeny is performed as described above.

The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

Applicant's or agent's file reference number A700/363	International application No. ...
--	-----------------------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 17, line 31	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution Agricultural Research Culture Collection (NRRL)	
Address of depositary institution (including postal code and country) 1815 North University Street Peoria, Illinois 61604 US	
Date of deposit January 12, 1996	Accession Number NRRL B-21517
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
1 microorganism is disclosed in the specification. In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication or the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28(4) EPC).	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<div style="border: 1px solid black; padding: 5px;"> <p>For receiving Office use only</p> <p><input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer <i>Therese Lind</i></p> </div>	<div style="border: 1px solid black; padding: 5px;"> <p>For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p> </div>

WHAT IS CLAIMED IS:

1. A method of altering glutenin content in the endosperm of a wheat plant, the method comprising:

stably transforming a parental wheat plant with a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and a seed-specific promoter to produce a stably transformed wheat plant; and

selecting a progeny of said stably transformed wheat plant having a glutenin content in the endosperm of a mature seed at least about 30% different than the glutenin content in the endosperm of a mature seed from the parental wheat plant.

2. The method of claim 1, wherein the nucleic acid is stably transformed into the first wheat plant using particle bombardment.

3. The method of claim 1, wherein the nucleic acid encodes a chimeric glutenin polypeptide.

4. The method of claim 1, wherein the chimeric polypeptide comprises sequences from an x-type glutenin polypeptide and a y-type glutenin polypeptide.

5. The method of claim 1, wherein the x-type polypeptide is encoded by the Glu-D1-1b gene.

6. The method of claim 1, wherein the y-type polypeptide is encoded by the Glu-D1-2b gene.

7. The method of claim 1, wherein the promoter is from the Glu-D1-2b gene.

8. The method of claim 1, wherein the parental wheat plant is Bobwhite.

9. A wheat plant comprising a recombinant expression cassette comprising a nucleic acid encoding a glutenin polypeptide and a seed-specific promoter, the plant having a

glutenin content in the endosperm of a mature seed at least about 30% different than the glutenin content in the endosperm of a mature seed from a parental wheat plant.

10. The plant of claim 9, wherein the nucleic acid encodes a chimeric glutenin polypeptide.

11. The plant of claim 10, wherein the chimeric polypeptide comprises sequences from an x-type polypeptide and a y-type glutenin polypeptide.

12. The plant of claim 11, wherein the x-type polypeptide is encoded by the Glu-D1-1b gene.

13. The plant of claim 11, wherein the y-type polypeptide is encoded by the Glu-D1-2b gene.

14. The plant of claim 9, wherein the promoter is from the Glu-D1-2b gene.

15. The plant of claim 9, wherein the high molecular weight (HMW) glutenin accounts for at least about 15% of the total protein in the endosperm of the mature seed.

16. The plant of claim 9, wherein the parental wheat plant is Bobwhite.

17. A seed derived from the plant of claim 9.

18. A recombinant construct comprising a wheat glutenin gene promoter of about 400 to about 2800 nucleotides operably linked to a heterologous polynucleotide sequence encoding a chimeric glutenin polypeptide, wherein the chimeric glutenin polypeptide is encoded by polynucleotide sequences from the Glu-D1-1b gene and the Glu-D1-2b gene.

19. The recombinant construct of claim 18, wherein the wheat glutenin gene is Glu-D1-2b.

20. A plant comprising th recombinant construct of claim 18.

21. Wheat flour derived from th plant of claim 1.

1/3




FIG. 1

2/3




FIG. 2A





FIG. 2B

3/3

**FIG. 3**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/00510

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/200, 205, 250, DIG 58; 435/172.3, 240.4, 320.1; 536/23.6, 24.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CABA, MEDLINE, CAPLUS, GENBANK, BIOSIS

search terms: wheat, glutenin, transformation, Glu-D1-2b, Glu-D1-1b

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SHANI et al. Role of amino- and carboxy- terminal regions in the folding and oligomerization of wheat high molecular weight glutenin subunits. Plant Physiol. 1992, Vol. 98, pages 433-441, see entire document.	1-21
Y	WEEKS et al. Rapid production of multiple independent lines of fertile transgenic wheat (<i>Triticum aestivum</i>). Plant Physiol. 1993, Vol. 102, pages 1077-1084, especially pages 1078-1079.	1-21
Y	ANDERSON et al. Nucleotide sequences of the two high-molecular-weight glutenin genes from the D-genome of a hexaploid bread wheat, <i>Triticum aestivum</i> L. cv Cheyenne. Nucleic Acids Research. 1989, Vol. 17, No. 1, pages 461-462, see entire document.	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A document defining the general state of the art which is not considered to be of particular relevance	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E earlier document published on or after the international filing date	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &	document member of the same patent family
* O document referring to an oral disclosure, use, exhibition or other means		
* P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

20 MARCH 1997

Date of mailing of the international search report

08 APR 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THOMAS HAAS

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/00510

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GALILI, G. Heterologous expression of a wheat high molecular weight glutenin gene in <i>Escherichia coli</i> . Proc. Natl. Acad. Sci. USA. October 1989, Vol. 86, pages 7756-7760, see entire document.	1-21

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/00510

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12N 15/00, 15/05, 15/09, 15/11, 15/29, 15/64, 15/82; A01H 1/00, 3/00, 4/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

800/200, 205, 250, DIG 58; 435/172.3, 240.4, 320.1; 536/23.6, 24.1





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00, 15/05, 15/09, 15/11, 15/29, 15/64, 15/82, A01H 1/00, 3/00, 4/00		A1	(11) International Publication Number: WO 97/25419
(21) International Application Number: PCT/US97/00510		(43) International Publication Date: 17 July 1997 (17.07.97)	
(22) International Filing Date: 10 January 1997 (10.01.97)		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 08/586,331 16 January 1996 (16.01.96) US			
(71) Applicant: THE UNITED STATES OF AMERICA , represented by TH [US/US]; E SECRETARY OF AGRICULTURE , Washington, DC 20250 (US).			
(72) Inventors: BLECHL, Ann, E. ; 1005 Buchanan Street, Albany, CA 94706 (US). ANDERSON, Olin, D. ; 208 Poshard Street, Pleasant Hill, CA 94523 (US).			
(74) Agents: PENDORF, Stephan, A. et al. ; Suite 1000, 600 N. Westshore Boulevard, Tampa, FL 33609 (US).			

Published
With international search report.

(54) Title: **GLUTENIN GENES AND THEIR USES****Y-type HMW-glutenin****X-type HMW-glutenin**

(57) Abstract

The present invention is directed to methods of improving flour by altering glutenin content in seeds of plants, particularly wheat. The invention relates to methods of introducing a recombinant construct comprising a glutenin gene into a parental plant.

* (Referred to in PCT Gazette No. 43/1997, Section II)

** (Referred to in PCT Gazette No. 49/1997, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam